

⑫

EUROPEAN PATENT APPLICATION

⑲ Application number: 88308916.1

⑤① Int. Cl.⁴: **C 07 K 7/00**
A 61 K 37/02

⑳ Date of filing: 26.09.88

③① Priority: 24.09.87 US 100571

④③ Date of publication of application:
29.03.89 Bulletin 89/13

⑥④ Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

⑦① Applicant: The Administrators of The Tulane Educational Fund
1430 Tulane Avenue
New Orleans Louisiana 70012 (US)

⑦② Inventor: Coy, David H.
4319 Perrier Street
New Orleans Louisiana 70115 (US)

Moreau, Jacques-Pierre
159 Westboro Road
Upton Massachusetts 05168 (US)

⑦④ Representative: Deans, Michael John Percy et al
Lloyd Wise, Tregear & CO. Norman House 105-109 Strand
London WC2R OAE (GB)

The microorganism(s) has (have) been deposited with American Type Culture Collection under number(s) CRL-1492.

⑤④ Therapeutic peptides.

⑤⑦ A linear peptide which is an analog of a naturally occurring, biologically active bombesin having an active site and a binding site responsible for binding of bombesin to a receptor on a target cell, cleavage of a peptide bond in the active site of the naturally occurring peptide being unnecessary for in vivo biological activity, the analog having a non-peptide bond instead of a peptide bond between an amino acid of the active site and an adjacent amino acid, and having the same binding site as the naturally occurring peptide, so that the analog, is capable of acting as a competitive inhibitor of naturally occurring bombesin by binding to the receptor and, by virtue of the non-peptide bond, failing to exhibit the in vivo activity of naturally occurring bombesin.

EP 0 309 297 A2

Description

Therapeutic Peptides



Background of the Invention

This invention relates to therapeutic peptides useful, e.g., in cancer therapy.

The amphibian peptide bombesin, pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-Leu-Met-NH₂ (Anastasi et al., *Experientia* 27:166-167 (1971)), is closely related to the mammalian gastrin-releasing peptides (GRP), e.g., the porcine GRP, H₂N-Ala-Pro-Val-Ser-Val-Gly-Gly-Gly-Thr-Val-Leu-Ala-Lys-Met-Tyr-Pro-Arg-Gly-Asn-His-Trp-Ala-Val-Gly-His-Leu-Met-(NH₂) (McDonald et al., *Biochem. Biophys. Res. Commun.* 90:227-233 (1979)) and human GRP, H₂N-Val-Pro-Leu-Pro-Ala-Gly-Gly-Gly-Thr-Val-Leu-Thr-Lys-Met-Tyr-Pro-Arg-Gly-Asn-His-Trp-Ala-Val-Gly-His-Leu-Met-NH₂. Bombesin has been found to be an autocrine or paracrine mitotic factor for a number of human cancer cell lines, including small-cell lung carcinoma (SCLC) (Haveman et al., eds. *Recent Results in Cancer Research - Peptide Hormones in Lung Cancer*, Springer-Verlag, New York:1986). A number of these cancers are known to secrete peptide hormones related to GRP or bombesin.

Consequently, antagonists to bombesin have been proposed as agents for the treatment of these cancers. Cuttitta et al. demonstrated that a specific monoclonal antibody to bombesin inhibited *in vivo* the growth of a human small-cell lung cancer cell line xenografted to nude mice (Cuttitta et al., *Cancer Survey* 4:707-727 (1985)). In 3T3 murine fibroblasts which are responsive to the mitotic effect of bombesin, Zachary and Rozengurt observed that a substance P antagonist (Spantide) acted as a bombesin antagonist (Zachary et al., *Proc. Natl. Acad. Sci. (USA)*, 82:7616-7620 (1985)). Heinz-Erian et al. replaced His at position 12 in bombesin with D-Phe and observed bombesin antagonist activity in dispersed acini from guinea pig pancreas (Heinz-Erian et al., *Am. J. of Physiol.* 252:G439-G442 (1987)). Rivier reported on work directed toward restricting the conformational freedom of the bioactive C-terminal decapeptide of bombesin by incorporating intramolecular disulfide bridges; however, Rivier mentioned that, so far, bombesin analogs with this modification fail to exhibit any antagonist activity (Rivier et al., "Competitive Antagonists of Peptide Hormones," in Abstracts of the International Symposium on Bombesin-Like Peptides in Health and Disease, Rome (October, 1987).

Abbreviations (uncommon):



Pal = 3-pyridyl-alanine

Nal = naphthylalanine

Summary of the Invention

In general, the invention features a linear (i.e., non-cyclic) peptide which is an analog of a naturally occurring, biologically active bombesin having an active site and a binding site responsible for the binding of bombesin to a receptor on a target cell.

the analog having a non-peptide bond instead of a peptide bond between an amino acid of the active site and an adjacent amino acid, the analog being capable of binding to the receptor, so that the analog is capable of acting as a competitive inhibitor of naturally occurring bombesin by binding to the receptor and, by virtue of the non-peptide bond, failing to exhibit the *in vivo* activity of naturally occurring bombesin. (A detailed discussion of the chemistry of non-peptide bonds is given in Coy et al. (1988) *Tetrahedron* 44,3:835-841, hereby incorporated by reference.)


Preferably, naturally occurring bombesin is characterized in that one or more amino acids in the amino terminal half of bombesin are hydrogen bonded to one or more amino acids in the carboxy terminal half of bombesin, and the non-peptide bond of the linear peptide decreases that hydrogen bonding, thereby destroying biological activity. It is believed that many of the linear peptides of the invention are analogs of bombesin whose biological activity depends at least in part on their ability to form tertiary "hairpin"

configurations in which amino acids in the amino terminal ("left") half of the molecule are hydrogen bonded to amino acids in the carboxy terminal ("right") half of the molecule, and that the pseudopeptide bond introduced according to the invention interferes with this hydrogen bonding, hindering the formation of the hairpin configuration on which activity depends. One may expect the loss of the ability to hydrogen bond to affect the biological activity of the molecule either by the loss of structural stability conferred by the transannular bonding or by the inability of the backbone to hydrogen bond to the receptor. Additionally, the increased flexibility of the molecule about the reduced bond compared with the rigidity of the normal peptide amide bond is expected to alter the conformational integrity of the molecule and thus its biological activity.

It is apparent from the above that the linear peptides for which introduction of a pseudopeptide bond is useful in creating or enhancing antagonist activity are those in which activity is associated with a site within the amino acid chain.

The pseudopeptide bond can be introduced in a region involved in receptor binding, or in a non-binding region; it has been shown (Nagain et al., *Peptides*, 8:1023-28 (1987)) that a pseudopeptide bond introduced in the binding region does not prevent binding. Generally, useful classes of peptides in which this modification can be made are those in which at least one amino acid involved in the active site is located in the carboxy terminal half of the molecule; the non-peptide bond is introduced between this amino acid and one adjacent to it.

One class of peptides of the invention is an effective bombesin antagonist peptide of formula (1):



wherein

- A¹ = pGlu or is deleted;
- A² = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, β -naphthylalanine or is deleted;
- A³ = Arg, D-Arg, Lys, D-Lys or is deleted;
- A⁴ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, β -naphthylalanine or is deleted;
- A⁵ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, D-Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, β -naphthylalanine D-Ala or is deleted;
- A⁶ = Gln, Asn, Gly, Ala, D-Ala, N-Ac-D-Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, p-Glu, β -naphthylalanine or is deleted;
- A⁷ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, D-Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, His, or β -naphthylalanine;
- A⁸ = Trp;
- A⁹ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, or β -naphthylalanine;
- A¹⁰ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, or β -naphthylalanine;
- A¹¹ = Gly, or D-Ala;
- A¹² = His, Phe, or p-X-Phe (X = F, Cl, Br, OH, CH₃);
- A¹³ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, β -naphthylalanine;
- A¹⁴ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, or β -naphthylalanine;


provided that

each R₁, R₂, R₃ and R₄ independently, is H, C₁₋₁₂ alkyl, C₇₋₁₀ phenylalkyl, COE₁ (where E₁ is C₁₋₂₀ alkyl, C₃₋₂₀ alkenyl, C₃₋₂₀ alkynyl, phenyl, naphthyl, or C₇₋₁₀ phenylalkyl), or COOE₂ (where E₂ is C₁₋₁₀ alkyl or C₇₋₁₀ phenylalkyl), and R₁ and R₂ are bonded to the N-terminal amino acid of said peptide, which can be A¹, A², A³, A⁴, A⁵, A⁶, or A⁷, provided that when one of R₁ or R₂ is COE₁ or COOE₂, the other must be H, and when one of R₃ or R₄ is COE₁ or COOE₂ the other must be H, and further provided that when A¹ = pGlu, R₁ must be H and R₂ must be the portion of Glu that forms the imine ring in pGlu; and for each of the residues A⁷, A⁸, A⁹, A¹¹, A¹², and A¹³, independently, the carbon atom participating in the amide bond between that residue and the

nitrogen atom of the alpha amino group of the adjacent amino acid residue may be a carbonyl carbon or may be reduced to a methylene carbon, provided that at least one such carbon atom must be reduced to a methylene carbon (i.e., at least one of the subject peptide CONH bonds must be replaced by a non-peptide, i.e., pseudopeptide, CH₂NH bond); or a pharmaceutically acceptable salt thereof. (Where no D- or L-isomeric designation is given herein, the naturally occurring L-isomer is intended.)

Preferably, an effective bombesin antagonist peptide has, for each of the residues A¹¹, A¹², and A¹³, independently, the carbon atom participating in the amide bond between that residue and the nitrogen atom of the alpha amino group of the adjacent amino acid residue which may be a carbonyl carbon or may be reduced to a methylene carbon, provided that at least one such carbon atom must be reduced to a methylene carbon; or a pharmaceutically acceptable salt thereof. Most preferably, the bombesin antagonist peptide has A¹ through A⁶ deleted and the carbon atom participating in the amide bond between Leu¹³ and Leu¹⁴ is a methylene carbon, or a pharmaceutically acceptable salt thereof

Another class of peptides of the invention are bombesin-related antagonist peptides derived from litorin and of the amino acid formula




wherein A¹ is pGlu; A² is Gln; A³ is Trp; A⁴ is Ala; A⁵ is Val; A⁶ is Gly or D-Ala; A⁷ is His; A⁸ is Phe or Leu; and A⁹ is Met or Leu; provided that the carbon atom participating in the amide bond between the A⁶ residue and the nitrogen atom of the alpha amino group of the adjacent amino acid residue may be a carbonyl carbon or may be reduced to a methylene carbon, or a pharmaceutically acceptable salt thereof.


Peptides of the invention that contain a pseudopeptide bond substitution within the active site of the naturally occurring peptide are antagonists to the biological activity of the naturally occurring bombesin peptide, with one exception which we have observed; the linear analog of bombesin BIM-26027 [Val¹⁰ψ[CH₂NH]Leu¹⁴]BN is an agonist of the biological activity of naturally occurring bombesin. (Non-peptide bonds are symbolized herein by "ψ[CH₂NH]" or "ψ".) Therefore, a third class of peptides of the invention are effective bombesin agonists of the formula (1) recited above, including, for the residue A¹⁰, the carbon atom participating in the amide bond between that residue and the nitrogen atom of the alpha amino group of the adjacent amino acid residue may be a carbonyl carbon or may be a non-peptide bond, provided that the non-peptide bond may be a carbonyl carbon having been reduced to a methylene carbon; further provided that at least one such carbon atom must be reduced to a methylene carbon; or a pharmaceutically acceptable salt thereof. Most preferred is the bombesin agonist having the formula pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-Leu-Leu[Val¹⁰ψ[CH₂NH]Leu¹⁴].

Other agonist analogues are peptides in which either the pseudopeptide bond is not located in the active site of the naturally occurring peptide, or in which two amino acid residues of the active site are replaced by statine or AHPPA.

(Statine has the chemical structure



and statine-amide has the structure



and AHPPA has the formula:

(3S,4S)-4-amino-3-hydroxy-5-phenylpentanoic acid.) Therefore, a fourth class of peptides of the invention is an effective bombesin agonist which is an analog of naturally occurring, biologically active bombesin having an active site, which includes positions A⁹, A¹⁰, A¹¹, A¹², A¹³, and A¹⁴, and a binding site responsible for the binding of bombesin to a receptor on a target cell, the analog having either (a) a non-peptide bond outside of the active site of bombesin, or (b) having at least one statine or AHPPA residue in place of two naturally occurring amino acids of the active site; and further, the peptide can contain statine or AHPPA when all bonds between amino acid residues are peptide bonds and, further, when an amino acid residue is statine or AHPPA, the amino acid to the right of it in the formula is deleted, so that the analog is capable of binding to the receptor and, by virtue of the statine or AHPPA residue, exhibiting enhanced *in vivo* biological activity compared to naturally occurring bombesin. Most preferred in this class is the bombesin agonist having the amino acid formula pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-[Sta¹³,Des Met¹⁴].

The bombesin antagonists and agonists of the invention are suitable for the treatment of all forms of cancer where bombesin-related substances act as autocrine or paracrine mitotic factors, especially pancreas and small-cell lung carcinoma

In formula (1), when R₁, R₂, R₃ or R₄ is an aromatic, lipophilic group, the *in vivo* activity can be long lasting, and delivery of the compounds of the invention to the target tissue (e.g., the lungs) can be facilitated

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Description of the Preferred Embodiments

We will first briefly describe the table.

Table

Table I shows formulas for the pseudo-peptide analogues and results of *in vitro* inhibition of [¹²⁵I] GRP binding to cerebral cortex and 3T3 fibroblast or murine fibroblast bombesin receptors, and bombesin-stimulated [³H]Thymidine uptake by cultured 3T3 cells.

We now describe the structure, synthesis, and use of the preferred embodiments of the invention.

Structure

The peptides of the invention all have a non-peptide bond in at least one of the indicated position, except for the statine or AHPPA substituted analogs, such as sta¹³-des Met¹⁴ bombesin. By non-peptide bond is meant that the carbon atom participating in the bond between two residues is reduced from a carbonyl carbon to a methylene carbon. The peptide bond reduction method which yields this non-peptide bond is described in Coy et al., U.S. patent application, Serial No. 879,348, assigned to the same assignee as the present application, hereby incorporated by reference. Any one or all of the amino acids in positions 1 through 6 of the bombesin antagonists may be deleted from the peptides, and the peptides are still active as antagonists or agonists.

The peptides of the invention can be provided in the form of pharmaceutically acceptable salts. Examples of preferred salts are those with therapeutically acceptable organic acids, e.g., acetic, lactic, maleic, citric, malic, ascorbic, succinic, benzoic, salicylic, methanesulfonic, toluenesulfonic, or pamoic acid, as well as polymeric acids such as tannic acid or carboxymethyl cellulose, and salts with inorganic acids such as the hydrohalic acids, e.g., hydrochloric acid, sulfuric acid, or phosphoric acid.

Synthesis of Bombesin Antagonists

The synthesis of the bombesin antagonist pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-Leuψ[CH₂-NH]Leu-NH₂ follows. Other bombesin antagonists and agonists can be prepared by making appropriate modifications of the following synthetic method.

The first step is the preparation of the intermediate pGlu-Gln-Arg(tosyl)-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His(benzyloxycarbonyl)-Leuψ[CH₂NH] Leu-benzhydrylamine resin, as follows.

Benzhydrylamine-polystyrene resin (Vega Biochemicals, Inc.) (0.97 g, 0.5 mmole) in the chloride ion form is placed in the reaction vessel of a Beckman 990B peptide synthesizer programmed to perform the following reaction cycle: (a) methylene chloride; (b) 33% trifluoroacetic acid (TFA) in methylene chloride (2 times for 1 and 25 min. each); (c) methylene chloride; (d) ethanol; (e) methylene chloride; and (f) 10% triethylamine in chloroform.

The neutralized resin is stirred with alpha-t-butoxycarbonyl(Boc)-leucine and diisopropylcarbodiimide (1.5 mmole each) in methylene chloride for 1 hour, and the resulting amino acid resin is then cycled through steps (a) to (f) in the above wash program Boc leucine aldehyde (1.25 mmoles), prepared by the method of Fehrentz and Castro, Synthesis, p. 676 (1983), is dissolved in 5 ml of dry dimethylformamide (DMF) and added to the resin TFA salt suspension followed by the addition of 100 mg (2 mmoles) of sodium cyanoborohydride (Sasaki and Coy, Peptides 8:119-121 (1987); Coy et al., *Id.*). After stirring for 1 hour, the resin mixture is found to be negative to ninhydrin reaction (1 min.), indicating complete derivatization of the free amino group.

The following amino acids (1.5 mmole) are then coupled successively in the presence diisopropylcarbodiimide (1.5 mmole), and the resulting amino acid resin is cycled through washing/deblocking steps (a) to (f) in the same procedure as above: Boc-His(benzylloxycarbonyl), Boc-Gly, Boc-Val, Boc-Ala, Boc-Trp, Boc-Gln (coupled in the presence of equivalent of hydroxybenzotriazole), Boc-Asn (coupled in the presence of 1 equivalent of hydroxybenzotriazole), Boc-Gly (coupled as a 6 M excess of the p-nitrophenyl ester), Boc-Leu, Boc-Arg(tosyl), Boc-Gln (coupled as a 6 M excess of the p-nitrophenylester), and pGlu. The completed resin is then washed with methanol and air dried.

The resin described above (1.6 g, 0.5 mmole) is mixed with anisole (5 ml) and anhydrous hydrogen fluoride (35 ml) at 0°C and stirred for 45 min. Excess hydrogen fluoride is evaporated rapidly under a stream of dry nitrogen, and free peptide is precipitated and washed with ether. The crude peptide is dissolved in a minimum volume of 2 M acetic acid and eluted on a column (2.5 x 100 mm) of Sephadex G-25 (Pharmacia Fine Chemicals, Inc.). Fractions containing a major component by uv absorption and thin layer chromatography (TLC) are then pooled, evaporated to a small volume and applied to a column (2.5 x 50 cm) of octadecylsilane-silica (Whatman LRP-1, 15-20 µm mesh size).

The peptide is eluted with a linear gradient of 0-30% acetonitrile in 0.1% trifluoroacetic acid in water. Fractions are examined by TLC and analytical high performance liquid chromatography (HPLC) and pooled to give maximum purity. Repeated lyophilization of the solution from water gives 60 mg of the product as a white, fluffy powder.

The product is found to be homogeneous by HPLC and TLC. Amino acid analysis of an acid hydrolysate confirms the composition of the peptide. The presence of the Leuψ[CH₂-NH]Leu bond is demonstrated by fast atom bombardment mass spectrometry.

pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Alaψ[CH₂-NH]Val-Gly-His-Leu-Met-NH₂ and pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-Leuψ[CH₂NH]Met-NH₂ are prepared in similar yields in an analogous fashion by appropriately modifying the above procedure

A statine or AHPPA residue can be substituted in place of any two amino acids of the peptide, where the peptide contains no pseudopeptide bonds. For example, sta¹³-des Met¹⁴ bombesin was prepared in an analogous fashion by first coupling statine to the resin and then proceeding with the addition of Boc-His(benzylcarbonyl). Statine or Boc-statine can be synthesized according to the method of Rich et al., 1978, J. Organic Chem. 43: 3624; and Rich et al., 1980, J. Med. Chem. 23: 27, and AHPPA can be synthesized according to the method of Hui et al., 1987, J. Med. Chem. 30: 1287.

Synthesis of Sta¹³-Des-Met¹⁴ Bombesin

Solid-phase synthesis of the peptide pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala Val-Gly-His-Sta-NH₂ was accomplished through the use of the following procedures in which alpha-t-butoxycarbonyl statine (prepared by the procedure of Rich et al., J. Org. Chem. 1978, 43, 3624) is first coupled to methylbenzhydrylamine-polystyrene resin. After acetylation, the intermediate p-Glu-Gln-Arg(tosyl)-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His(benzylloxycarbonyl)-Sta-methylbenzhydrylamine resin is prepared. The synthetic procedure used for this preparation follows in detail:

1. Incorporation of alpha-t-butoxycarbonyl statine on methylbenzhydrylamine resin.

Methylbenzhydrylamine-polystyrene resin (Vega Biochemicals, Inc.) (1.0 g, 0.73 mmol) in the chloride ion form is placed in the reaction vessel of a Vega 250C Coupler peptide synthesizer. The synthesizer was programmed to perform the following reactions: (a) methylene chloride; (b) 10% triethylamine in chloroform; (c) methylene chloride; and (d) dimethylformamide.

The neutralized resin is mixed for 18 hours with the preformed active ester made from alpha-t-butoxycarbonyl statine (1.46 mmol), diisopropyl carbodiimide (2 mmol), and hydroxybenzotriazole hydrate (1.46 mmol in dimethylformamide at 0° C. for one hour. The resulting amino acid resin is washed on the synthesizer with dimethylformamide and then methylene chloride. The resin mixture at this point was found by the Kaiser ninhydrin test (5 minutes) to have an 84% level of statine incorporation on the resin.

Acetylation was performed by mixing the amino-acid resin for 15 minutes with N-acetyl imidazole (5 mmol) in methylene chloride. Derivatization to the 94-99% level of the free amino groups of the resin was indicated by the Kaiser ninhydrin test (5 minutes). The Boc-statine-resin is then washed with methylene chloride.

2. Couplings of the Remaining Amino Acids.

The peptide synthesizer is programmed to perform the following reaction cycle: (a) methylene chloride; (b) 33% trifluoroacetic acid (TFA) in methylene chloride (2 times for 5 and 25 min. each); (c) methylene chloride; (d) isopropyl alcohol; (e) 10% triethylamine in chloroform; and (f) methylene chloride.

The following amino acids (2.19 mmol) are then coupled successively by diisopropyl carbodiimide (4 mmol) alone or diisopropyl carbodiimide (4 mmol) plus hydroxybenzotriazole hydrate (1.47 or 0.73 mmol) and the resulting peptide-resin is washed on the synthesizer with dimethylformamide and then methylene chloride, and then cycled through the washing and deblocking steps (a) to (f) in the procedure described above.

Boc-His (benzylloxycarbonyl) (coupled in the presence of 2 equivalents hydroxybenzotriazole); Boc-Gly;

Boc-Val; Boc-Ala; Boc-Trp; Boc-Gln and Boc Asn (coupled as the preformed hydroxybenzotriazole active esters made by reaction at 0° C. for one hour with 1 equivalent hydroxybenzotriazole hydrate); Boc-Gly; Boc-Leu; Boc-Arg(tosyl), Boc-Gln, and pGlu (also coupled as the preformed active esters of hydroxybenzotriazole made by reaction at 0° C. for one hour with 1 equivalent hydroxybenzotriazole hydrate). The completed peptide-resin is then washed with methanol and air dried.

The peptide-resin described above (1.60 g, 0.73 mmol) is mixed with anisole (2.5 mL), dithioerythritol (50 mg), and anhydrous hydrogen fluoride (30 mL) at 0° C. for one hour. Excess hydrogen fluoride is evaporated rapidly under a stream of dry nitrogen, and the free peptide is precipitated and washed with ether. The crude peptide is dissolved in 100 mL of 1 M acetic acid and the solution is then evaporated under reduced pressure. The crude peptide is dissolved in a minimum volume of methanol/water 1/1 and triturated with 10 volumes of ethyl acetate.

The triturated peptide is applied to a column (9.4 mm I.D. x 50 cm) of octadecylsilane-silica (Whatman Partisil 10 ODS-2 M 9). The peptide is eluted with a linear gradient of 20-80% of 20/80 0.1% trifluoroacetic acid/acetoneitrile in 0.1% trifluoroacetic acid in water. Fractions are examined by TLC and analytical high performance liquid chromatography (HPLC) and pooled to give maximum purity. Lyophilization of the solution from water gives 77 mg of the product as a white fluffy powder.

Other compounds can be prepared as above and tested for effectiveness as agonists or antagonists in the following test program.

Phase 1 - 3T3 Peptide Stimulated [³H] Thymidine Uptake Assay

Cell Culture. Stock cultures of Swiss 3T3 cells (American Type Culture Collection No. OCL 92) are grown in Dulbecco's Modified Eagles Medium (DMEM) supplemented with 10% fetal calf serum in humidified atmosphere of 10% CO₂/90% air at 37° C. For experimental use, the cells are seeded into 24-well cluster trays and used four days after the last change of medium. The cells are arrested in the G1/G0 phase of the cell cycle by changing to serum-free DMEM 24 hours prior to the thymidine uptake assay.

Assay of DNA Synthesis. The cells are washed twice with 1ml aliquots of DMEM (-serum) then incubated with DMEM (-serum), 0.5μM [methyl-³H] thymidine (20Ci/mmol, New England Nuclear), bombesin (1nM), and four concentrations of the test compounds (1, 10, 100, 1000nM) in a final volume of 0.5ml. After 28 hours at 37° C, [methyl-³H] thymidine incorporation into acid-insoluble pools is assayed as follows. The cells are washed twice with ice-cold 0.9% NaCl (1ml aliquots), and acid soluble radioactivity is removed by a 30 min. (4° C) incubation with 5% trichloroacetic acid (TCA). The cultures are then washed once (1ml) with 95% ethanol and solubilized by a 30 min. incubation (1ml) with 0.1N NaOH. The solubilized material is transferred to vials containing 15ml ScintA (Packard), and the radioactivity is determined by liquid scintillation spectrometry.

Phase 2 - Small Cell Carcinoma (SCLC) - Bombesin Stimulated [³H] Thymidine Uptake Assay

Cell Culture. Cultures of the human cell carcinoma cell line (NCI-H69) (obtained from the American Type Culture Association) are maintained in RPMI 1640 medium supplemented with 10% fetal calf serum in 10% CO₂/90% air at 37° C. Twenty-four hours prior to assay, the cells are washed with serum-free medium and seeded in 24-well cluster trays.

Assay of DNA Synthesis. Bombesin (1nM), 0.5μM [methyl-³H] thymidine (20 Ci/mmol, New England Nuclear), and four concentrations of the test compounds (1, 10, 100, 1000nM) are added to the cultures to achieve a final volume of 0.5 ml. After a 28 hr incubation at 37° C, the cells are collected onto GF/B glass fiber filters, and the DNA is precipitated with ice-cold TCA. [³H] thymidine incorporation into acid-insoluble fractions of DNA is determined by liquid scintillation spectrometry.

Phase 3 - Peptide-Induced Pancreatitis

Male, Sprague-Dawley rats (250g) are used for these experiments. The test compound, or 0.9% NaCl is administered s.c. 15 min. prior to the bombesin injection. Bombesin injections are given s.c. at a dose of 10 μg/kg, and blood samples are obtained at 1 hr. 30 min., 3hr. and 6hr. Plasma amylase concentration are determined by the Pantrak Amylase test.

Phase 4 - In Vitro Inhibition of [¹²⁵I] Gastrin Releasing Peptide (GRP) Binding to Bombesin Receptors

Membranes from various tissues (rat brain, rat pancreas, rat anterior pituitary, SCLC, 3T3 cells) are prepared by homogenization in 50mM TrisHCl containing 0.1% bovine serum albumin and 0.1mg/ml bacitracin followed by two centrifugations (39,000xg x 15 min., 4° C) with an intermediate resuspension in fresh buffer. For assay, aliquots (0.8ml) are incubated with 0.5nM [¹²⁵I]GRP (2000 Ci/mmol, Amersham Corp.) and various concentrations of the test compounds in a final volume of 0.5ml. After a 30 minute incubation at 4° C, the binding reaction is terminated by rapid filtration through Whatman GF/C filters that have been pre-soaked in 0.3% aqueous polyethyleneimine to reduce the level of nonspecific binding. The filters and tubes are washed three times with 4ml aliquots of ice-cold buffer, and the radioactivity trapped on the filters is counted by gamma-spectrometry. Specific binding is defined as the total [¹²⁵I]GRP bound minus that bound in the

presence of 1000nM bombesin.

Phase 5 - Inhibition of Gastrin Release

- 5 The stomachs of anesthetized rats are perfused with saline collected over 15 minute periods via pyloric cannulation while the test peptide is infused through the femoral vein for periods between 0 and 150 minutes.

Results of Tests of Test Peptides

- 10 A number of analogs of bombesin, each containing a non-peptide bond, were synthesized and tested in one or more of the above-described Phase 1 - 5 assays; the results of Phase 1, 2, and 4 tests are given in Table 1 attached hereto (analogs of bombesin are indicated by the symbol "BN"). Brain and 3T3 GRP receptor and thymidine uptake data are expressed in IC₅₀ (nM) Table 1 also gives results for non-peptide bond-containing analogs of three other naturally-occurring peptides, Substance P (which plays a role in the sensation of pain),
15 Neuromedin C, whose C-terminal seven amino acids are similar to those of bombesin, and litorin, whose eight C-terminal amino acids are identical to Bombesin, with the exception of a Phe substitution for Leu at position A¹³ of bombesin.

- In Table 1, the position of the non-peptide bond is indicated by the position of the symbol ψ ; i.e., ψ is always shown preceding the amino acid which, in that peptide, is bonded to the amino acid N-terminal to it via the non-peptide bond. Where no amino acid is specified under "structure", as in BIM-26034, the non-peptide bond links the two peptides represented by the numbers given as post-scripts (e.g., between amino acids 7 and 8 of BIM-26034, which otherwise is identical to naturally occurring bombesin).
20

- In Table 1, it can be seen that a preferred placement of the non-peptide bond in bombesin analogs is at the 13-14 position; two of the most active analogs (as indicated by a low GRP receptor IC₅₀ value) are BIM-26027 and BIM-26028. However, BIM-26027 causes proliferation of cancer cells (see Table 1, under thymidine uptake), and therefore is an agonist and not an antagonist. In general, compounds having the non-peptide bond at any position other than the active site of the peptide are agonists rather than antagonists. Table 1 also shows that when statine replaces the A¹³ and A¹⁴ residues of bombesin, the resultant bombesin analog BIM-26096 causes proliferation of cancer cells and is therefore an agonist. Bombesin superagonists may be
25 useful in cancer therapy, as suggested by Alexander et al., 1988, Cancer Research 48: 1439-1441, and Alexander et al., 1988, Pancreas 3:297-302, hereby incorporated by reference. Alexander et al. show that chronic bombesin treatment inhibited the growth of human ductal adenocarcinoma transplanted into athymic mice. These results were surprising for bombesin stimulates growth of normal pancreas tissue. The demonstration of both stimulatory and inhibitory activity suggests that bombesin interacts differently in normal and neoplastic pancreatic tissues.
30

- These observations prompted us to evaluate the affect of BIM-26096, a bombesin analogue which has bombesin-like agonist activity, on the *in vitro* growth of a pancreatic tumor cell line (AR42J, A.T.C.C. No. CRL1492. For these experiments, AR42J cells were subcultured into a 24-well culture plate in Dulbecco's modified Eagle's medium containing 10% fetal calf serum containing various concentrations (0.1-100nM) of BIM-26096. After a 36 hr incubation the cells were removed with a trypsin/EDTA solution and the number of
35 cells were determined using a Coulter Counter. The results are shown below:

45	<u>Treatment</u>	<u>Cell Count (% Control)</u>
	control	100
	BIM-26096 (0.1 nM)	78
50	BIM-26096 (1.0 nM)	73
	BIM-26096 (10 nM)	56
	BIM-26096 (100 nM)	52

- 55 These results indicate that the bombesin agonist, BIM-26096, has *in vitro* antiproliferative activity against the AR42J rat pancreas tumor.

- Finally, Table 1 also shows that bond placement, while important, is not the only factor influencing antagonist activity, and that amino acid substitutions at some positions exert influence as well; this is illustrated by BIM-26030, with Gly in position 11, which exhibited no antagonist activity. Table 1 also gives
60 negative results for analogs of Spantide ([D-Arg', D-Trp^{7,9} Leu"] Substance P, and Bombesin. Thus the non-peptide bond placement guidelines given herein should be used in conjunction with the routine assays described above to select useful antagonists or agonists.

- In a phase 5 assay, above, the results of which are not given in Table 1, BIM-26028 was shown to be a potent
65 inhibitor of bombesin - stimulated gastric acid secretion.

Use

The peptides of the invention may be administered to a mammal, particularly a human, in one of the traditional modes (e.g., orally, parenterally, transdermally, or transmucosally), in a sustained release formulation using a biodegradable biocompatible polymer, or by on-site delivery (e.g., in the case of anti-cancer bombesin to the lungs) using micelles, gels and liposomes.

The bombesin antagonists and agonists of the invention are suitable for the treatment of all forms of cancer where bombesin-related substances act as autocrine or paracrine mitotic agents, particularly small-cell lung carcinoma. The peptides can also be used for the inhibition of gastric acid secretion, the symptomatic relief and/or treatment of exocrine pancreatic adenocarcinoma, and the restoration of appetite to cachexic patients. The peptides can be administered to a human patient in a dosage of 0.5 µg/kg/day to 5 mg/kg/day. For some forms of cancer, e.g., small cell lung carcinoma, the preferred dosage for curative treatment is 250mg/patient/day.

5

10

15

20

25

30

35

40

45

50

55

60

65

Table 1

<u>Code</u>	<u>Structure</u>	<u>Brain GRP Receptor IC50(nM)</u>	<u>3T3 GRP Receptor IC50(nM)</u>	<u>Thym. Uptake IC50(nM)</u>
BIM-26025	[His ¹² ψ[CH ₂ NH]Leu ¹⁴]BN	>1000		
BIM-26026	[Ala ⁹ ψ[CH ₂ NH]Leu ¹⁴]BN	>1000		1574
BIM-26027	[Val ¹⁰ ψ[CH ₂ NH]Leu ¹⁴]BN	0.48	2.3	agonist EC50=0.0
M BIM-26028	[Leu ¹³ ψ[CH ₂ NH]Leu ¹⁴]BN	13		14.7
BIM-26030	[Gly ¹¹ ψ[CH ₂ NH]Leu ¹⁴]BN	>1000		
BIM-26034	[Gln ⁷ ψ[CH ₂ NH]]BN	>1000		
BIM-26036	[Des-pGlu ¹ ,Gln ² , Ala ⁹ ψ[CH ₂ NHPhe ¹²]BN	>1000		
BIM-26046	[Gly ¹¹ ψ[CH ₂ NH]D-Phe ¹² , Leu ¹⁴]BN	>1000		
BIM-26048	[D-Phe ¹² ψ[CH ₂ NH]Leu ¹³ , Leu ¹⁴]BN	>1000		
BIM-26056	[Leu ¹⁰ ψ[CH ₂ NH] Leu ¹¹ NH ₂]Substance P	>1000		
BIM-26057	[Cys ⁹ ,ψLeu ¹³ ,Cys ¹⁴]BN	>1000		

<u>Code</u>	<u>Structure</u>	Brain GRP Receptor <u>IC50(nM)</u>	3T3 GRP Receptor <u>IC50(nM)</u>	Thym. Uptake <u>IC50(nM)</u>
BIM-26061	[D-pGlu, D-Ala ⁵ , Ψ Leu ⁷ Met ⁸]BN	>1000		
BIM-26062	[Phe ¹³ Ψ [CH ₂ NH]Leu ¹⁴]BN	>1000		437
BIM-26063	[des-Glu ⁷ , Leu ¹³ Ψ [CH ₂ NH]Leu ¹⁴]BN		>1000	
BIM-26064	[Ψ Leu ¹⁰ , Nle ¹¹]Spantide	>1000		
BIM-26067	[des-Gln ⁷ , Ψ Leu ¹³⁻¹⁴]BN	>1000		
BIM-26068	[Ψ Leu ¹³ , Phe ¹⁴]BN	2.9		70
BIM-26070	[Ψ D-Trp ⁹ , Nle ¹¹]Spantide	>1000		
BIM-26071	[Tyr ⁴ , Ψ Leu ¹³ [CH ₂ NH]-Met ¹⁴]BN	34	16	104
BIM-26072	[Cys ⁹ , Leu ¹³ [CH ₂ NH] Cys ¹⁴]BN	>1000		
BIM-26073	[Cys ⁹ , Ψ Leu ¹³ [CH ₂ NH] Cys ¹⁴]BN	>1000		
BIM-26074	[Des-Gln ⁷ , Ψ Leu ¹³ [CH ₂ NH] Leu ¹⁴]BN	>1000		
BIM-26075	[D-Phe ¹¹ , Leu ¹³ Ψ [CH ₂ NH] Leu ¹⁴]BN	>1000		

<u>Code</u>	<u>Structure</u>	Brain	3T3 GRP	Thym.
		GRP Receptor <u>IC50(nM)</u>	Receptor <u>IC50(nM)</u>	Uptake <u>IC50(nM)</u>
BIM-26076	[D-Phe ¹¹ , ψ Leu ¹³⁻¹⁴]BN	>1000		
BIM-26077	[D-Ala ⁵ , ψ Leu ¹³⁻¹⁴]Leu ¹⁴ BN	517	196	1001
BIM-26078	[D-Ala ¹¹ , ψ Leu ¹³⁻¹⁴] Leu ¹⁴ BN	>1000		70
BIM-26079	[ψ Phe ⁷ , Leu ¹¹]Spantide	>1000		
BIM-26080	[ψ Gln ⁶ -Nle ¹¹]Spantide	>1000		
BIM-26081	[ψ D-Trp ⁷ -Nle ¹¹]Spantide	>1000		
BIM-26082	[ψ Phe ⁸ -Nle ¹¹]Spantide	>1000		
BIM-26083	[ψ Gln ⁶ -Nle ¹¹]Spantide	>1000		
BIM-26084	[ψ D-Trp ⁷ -Nle ¹¹]Spantide	>1000		
BIM-26085	[ψ Phe ⁸ -Nle ¹¹]Spantide	>1000		
BIM-26086	[D-Phe ¹² , ψ Leu[CH ₂ NH] Leu ¹⁴]BN	>1000		
BIM-26088	[ψ Gly ⁹ [CH ₂ NH]Leu ¹⁴] Spantide	>1000		
BIM-26089	[ψ Gln ⁶ [CH ₂ NH]Leu ¹¹] Spantide	>1000		
BIM-26090	[γ Phe ⁷ , Leu ¹¹]Substance P			>1000

<u>Code</u>	<u>Structure</u>	Brain	3T3 GRP	Thym.
		GRP Receptor <u>IC50(nM)</u>	Receptor <u>IC50(nM)</u>	Uptake <u>IC50(nM)</u>
BIM-26091	[ψ Phe ⁸ ,Leu ¹¹]Substance P			>1000
BIM-26092	[ψ Leu ⁹]Neuromedin C		242	466
BIM-26093	des ¹⁻⁴ ,D-Ala ⁵ ,His ⁷ ,Leu ¹³ ψ [CH ₂ NH]BN		82	171
BIM-26094	[D-Ala ^{5,11} ,Leu ¹³ ψ [CH ₂ NH] Leu ¹⁴]BN		1613	574
BIM-26095	[D-Ala ⁶ ,Leu ⁹ ψ [CH ₂ NH] Leu ¹⁰]Litorin		2623	1209
BIM-26096	[Sta ¹³ ,Des Met ¹⁴]BN	33		agonist EC50=3nM
BIM-26097	[Ac-Lys ⁷ , ψ Leu ¹³]BN ₇₋₁₄	1000		>1000
BIM-26098	[Lys ⁷ , ψ Leu ¹³]BN ₇₋₁₄	1000		
BIM-26099	[ψ Leu ¹³ ,Met]BN		73	78
BIM-26100	[Phe ⁸ ψ [CH ₂ NH]Leu ⁹]Litorin		74	22
BIM-26101	Leu ⁸ ψ [CH ₂ NH]Leu ⁹]Litorin		17.9	257
BIM-26102	ψ Phe ⁹ [CH ₂ NH]Met ¹⁰ NH ₂ Neuromedin B		184	>1000

5	Code	Structure	Brain		
			GRP Receptor IC50(nM)	3T3 GRP Receptor IC50(nM)	Thym. Uptake IC50(nM)
10	BIM-26103	Leu ¹³ ψ[CH ₂ NH]Met ¹⁴ NH ₂ A-Lytensin		>1000	>1000
15	BIM-26104	ψLeu ⁷ [CH ₂ NH]Met ⁸ NH ₂ GRP(20-27)			>1000
20	Spantide	[D-Arg ¹ ,D-Trp ^{7,9} ,Leu ¹¹] Substance P		3303	2171
25	Bombesin	pGlu-Gln-Arg-Leu-Gly-Asn- Gln-Trp-Ala-Val-Gly-His- Leu-Met-NH ₂	15	0.17	

40 Claims

1. A linear peptide which is an analog of naturally occurring, biologically active mammalian GRP or amphibian bombesin having an active site and a binding site responsible for the binding of bombesin to a receptor on a target cell, said analog having a non-peptide bond instead of a peptide bond between an amino acid of said active site and an adjacent amino acid, said analog being capable of binding to said receptor, so that said analog is capable of acting as a competitive inhibitor of said naturally occurring peptide by binding to said receptor and, by virtue of said non-peptide bond, failing to exhibit the in vivo activity of said naturally occurring bombesin.


2. The linear peptide of claim 1 wherein said naturally occurring bombesin is characterized in that one or more amino acids in the amino terminal half of bombesin are hydrogen bonded to one or more amino acids in the carboxy terminal half of bombesin, and said non-peptide bond of said linear peptide decreases said hydrogen bonding.

3. The linear peptide of claim 2 wherein said hydrogen bonded amino acids of said naturally occurring bombesin make up at least a portion of the active site of said naturally occurring bombesin, so that said active site is inactivated by the decrease in hydrogen bonding caused by said non-peptide bond.

4. A linear peptide which is an analog of naturally occurring, biologically active human bombesin which includes an active site comprising at least one amino acid in the carboxy terminal half of bombesin, said linear peptide including said amino acid in its carboxy terminal half, there being a non-peptide bond bonding said amino acid to an adjacent amino acid.

5. The linear peptide of claim 4 wherein said amino acid of said naturally occurring bombesin is hydrogen bonded to another, non-adjacent amino acid in said bombesin, and said non-peptide bond in said linear peptide causes a decrease in said hydrogen bonding which inactivates said bombesin.

6. An effective bombesin antagonistic peptide containing the amino acid formula:



wherein,

A¹ = pGlu, D or L, or is deleted.

A² = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, β -naphthylalanine or is deleted;

A³ = Arg, D-Arg, Lys, D-Lys or is deleted;

A⁴ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, β -naphthylalanine or is deleted;

A⁵ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, D-Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, β -naphthylalanine, D-Ala or is deleted;

A⁶ = Gln, Asn, Gly, Ala, D-Ala, N-Ac-D-Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, p-Glu, β -naphthylalanine or is deleted;

A⁷ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, D-Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, Lys, His, or α -naphthylalanine;

A⁸ = Trp, Met;

A⁹ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃) Trp, or β -naphthylalanine, D or L;

A¹⁰ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, Thr, or β -naphthylalanine;

A¹¹ = Gly, Phe, D or L;

A¹² = His, Phe, or p-X-Phe (X = F, Cl, Br, OH, CH₃) D or L;

A¹³ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, or β -naphthylalanine;

A¹⁴ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, or β -naphthylalanine;


provided that

each R₁, R₂, R₃, and R₄, independently, is H, C₁₋₁₂ alkyl C₇₋₁₀ phenylalkyl, COE₁ (where E₁ is C₁₋₂₀ alkyl, C₃₋₂₀ alkenyl, C₃₋₂₀ alkynyl, phenyl, naphthyl, or C₇₋₁₀ phenylalkyl), or COOE₂ (where E₂ is C₁₋₁₀ alkyl or C₇₋₁₀ phenylalkyl), and R₁ and R₂ are bonded to the N-terminal amino acid of said peptide, which can be A¹, A², A³, A⁴, A⁵, A⁶, or A⁷, and further provided that when one of R₁ or R₂ is COE₁ or COOE₂, the other must be H, and when one of R₃ or R₄ is COE₁ or COOE₂, the other must be H, and further provided that when A¹ = pGlu, R₁ must be H and R₂ must be the portion of Glu that forms the imine ring in pGlu; and for each of the residues A⁷, A⁸, A⁹, A¹¹, A¹², and A¹³, independently, the carbon atom participating in the amide bond between that residue and the nitrogen atom of the alpha amino group of the adjacent amino acid residue may be a carbonyl carbon or may be reduced to a methylene carbon, provided that at least one such carbon atom must be reduced to a methylene carbon; or a pharmaceutically acceptable salt thereof.

7. The effective bombesin antagonist peptide of claim 6 wherein A¹ through A⁶ are deleted and the carbon atom participating in the amide bond between Leu¹³ and Leu¹⁴ is a methylene carbon; or a pharmaceutically acceptable salt thereof.

8. The effective bombesin antagonist peptide of claim 6 wherein, for each of said residues A¹¹, A¹², and A¹³, independently, the carbon atom participating in the amide bond between that residue and the nitrogen atom of the alpha amino group of the adjacent amino acid residue may be a carbonyl carbon or may be reduced to a methylene carbon, provided that at least one such carbon atom must be reduced to a methylene carbon; or a pharmaceutically acceptable salt thereof.

9. An effective litorin antagonist peptide containing the amino acid formula:



wherein A¹ is A⁶ of Claim 6, A² is A⁷ of Claim 6, A³ is A⁸ of Claim 6, A⁴ is A⁹ of Claim 6, A⁵ is A¹⁰ of Claim 6, A⁶ is A¹¹ of Claim 6, A⁷ is A¹² of Claim 6, A⁸ is A¹³ of Claim 6, and A⁹ is A¹⁴ of Claim 6; provided that the carbon atom participating in the amide bond between the A⁸ residue and the nitrogen atom of the alpha amino group of the adjacent amino acid residue may be a carbonyl carbon or may be reduced to a methylene carbon; or a pharmaceutically acceptable salt thereof.

10. An effective bombesin agonist of the general formula of Claim 6 wherein, for the residue A¹⁰, the carbon atom participating in the amide bond between that residue and the nitrogen atom of the alpha amino group of the adjacent amino acid residue may be a carbonyl carbon or may be a non-peptide bond, provided that said non-peptide bond is said carbonyl carbon having been reduced to a methylene carbon, further provided that at least one such carbon atom must be reduced to a methylene carbon; or a pharmaceutically acceptable salt thereof.

11. A bombesin agonist having the amino acid formula


11. A bombesin agonist having the amino acid formula
pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Ψ [CH₂NH]-Gly-His-Leu-Leu.

12. An effective bombesin agonist having the amino acid formula of Claim 6 which is an analog of naturally occurring, biologically active mammalian GRP or amphibian bombesin having an active site, said active site includes the positions A⁹, A¹¹, A¹², A¹³, and A¹⁴ and a binding site responsible for the binding of said bombesin to a receptor on a target cell, said analog having either (a) said non peptide bond at residues other than within said active site, or (b) having at least one statine or AHPPA residue in place of two naturally occurring amino acids of said active site, and further provided that the peptide can contain statine or AHPPA when all bonds between amino acid residues are peptide bonds, and further provided that when an amino acid residue is statine or AHPPA, the amino acid to the right of it in the formula is deleted, so that said analog is capable of binding to said receptor and, by virtue of said statine or AHPPA residue, exhibiting enhanced in vivo biological activity compared to said naturally occurring bombesin.

13. A bombesin agonist having the amino acid formula

13. A bombesin agonist having the amino acid formula
pGlu-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-[Sta¹³,Des Met¹⁴].

14. The method of synthesizing an effective bombesin antagonistic peptide containing the following amino acids formula:



wherein, for the positions A¹ through A¹⁴,

A¹ = pGlu, D or L or is deleted;

A² = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, β -naphthylalanine or is deleted;

$A^3 = \text{Arg, D-Arg, Lys, D-Lys or is deleted;}$

A⁵ = Arg, D-Arg, Lys, D-Lys or is deleted,
A⁴ Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃),
Trp, B naphthylalanine or is deleted;

A⁵ = Gln, Asn, Gly, Ala, Leu, Ile, α-aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, B-naphthylalanine, D-Ala, or is deleted;

A⁶ = Gln, Asn, Gly, Ala, D-Ala, N-Ac-D-Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, D-Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, pGlu, B-naphthylalanine, or is deleted;

A⁷ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α-aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃). Trp, Lys, His, B-naphthylalanine;

$A^8 = \text{Trp or Met};$

A⁹ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α-aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, B-naphthylalanine, D or L;

A¹⁰ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α -aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or

CH₃), Trp, Thr, or B-naphthylalanine;

A¹¹ = Gly, Phe, D or L;

A¹² = His, Phe, or p-X-Phe (X = F, Cl, Br, OH or CH₃), D or L;

A¹³ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α-aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, B-naphthylalanine;

A¹⁴ = Gln, Asn, Gly, Ala, Leu, Ile, Nle, α-aminobutyric acid, Met, Val, Phe, p-X-Phe (X = F, Cl, Br, OH or CH₃), Trp, B-naphthylalanine;

provided that

each R₁, R₂, R₃, and R₄, independently, is H, C₁₋₁₂ alkyl, C₇₋₁₀ phenylalkyl, COE₁ (where E₁ is C₁₋₂₀ alkyl, C₃₋₂₀ alkenyl, C₃₋₂₀ alkynyl, [remainder of claim 6],

said method comprising the steps of (a) preparing benzyldrylamine-polystyrene resin, (b) neutralizing said resin and coupling a chosen amino acid for said A¹⁴ position to said resin, (c) coupling a chosen amino acid for said A¹³ position to said A¹⁴-resin, (d) derivatizing the free amino group of said A¹³-A¹⁴-resin, and (e) coupling successively to said derivatized amino acid resin a chosen amino acid for said A¹² through A¹ positions by repeating steps (d) and (e) for each of said amino acid positions.

15. The method of claim 14, wherein any or all of said positions A⁶ through A¹ are deleted.

16. A method for synthesizing an effective bombesin agonist of the general formula of claim 6 and according to the method of claim 6 wherein the carbon atom participating in the amide bond between said position A¹⁰ and the nitrogen atom of the alpha amino group of said position A¹¹ is a non-peptide bond, said non-peptide bond comprising a methylene carbon.

17. A method of synthesizing an effective bombesin agonist having the amino acid formula of claim 6 which is an analog of naturally occurring, biologically active mammalian GRP or amphibian bombesin, wherein said active site includes the positions A⁹, A¹¹, A¹², A¹³, and A¹⁴, and a binding site responsible for the binding of said bombesin to a receptor on a target cell, said method comprising either the steps of (a) preparing benzyldrylamine-polystyrene resin, (b) neutralizing said resin and coupling a chosen for said A¹⁴ amino acid position or alpha-t-butoxycarbonyl statine to said resin, (c) coupling a chosen amino acid for said A¹³ position or said statine for said A¹² position to said A¹⁴-resin, (d) derivatizing the free amino group of said A¹³-A¹⁴-resin or said statine-resin, and (e) coupling successively to said derivatized amino acid or statine resin a chosen amino acid for said A¹² through A¹ positions by repeating steps (d) and (e) for each of said amino acid positions; provided that when said statine is coupled to said resin, said statine-resin is acetylated before derivatization to free amino of groups; provided at least one statine or AHPPA residue replaces two naturally occurring amino acids of said active site; or the steps of the method of claim 6 wherein the non-peptide bond is outside of said active site.

THIS PAGE BLANK (USPTO)

⑫

EUROPEAN PATENT APPLICATION

⑲ Application number: 88308916.1

⑤① Int. Cl.⁵: **C07K 7/00, A61K 37/02,**
C07K 7/02

⑳ Date of filing: 26.09.88

③① Priority: 24.09.87 US 100571

④③ Date of publication of application:
29.03.89 Bulletin 89/13

⑧④ Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

⑧⑤ Date of deferred publication of the search report:
04.07.90 Bulletin 90/27

⑦① Applicant: **The Administrators of The Tulane Educational Fund**
1430 Tulane Avenue
New Orleans Louisiana 70012(US)

⑦② Inventor: **Coy, David H.**
4319 Perrier Street
New Orleans Louisiana 70115(US)
Inventor: **Moreau, Jacques-Pierre**
159 Westboro Road
Upton Massachusetts 05168(US)

⑦④ Representative: **Deans, Michael John Percy et al**
Lloyd Wise, Tregear & CO. Norman House
105-109 Strand
London WC2R OAE(GB)

⑤④ **Therapeutic peptides.**

⑤⑦ A linear peptide which is an analog of a naturally occurring, biologically active bombesin having an active site and a binding site responsible for binding of bombesin to a receptor on a target cell, cleavage of a peptide bond in the active site of the naturally occurring peptide being unnecessary for in vivo biological activity, the analog having a non-peptide bond instead of a peptide bond between an amino acid of the active site and an adjacent amino acid, and having the same binding site as the naturally occurring peptide, so that the analog, is capable of acting as a competitive inhibitor of naturally occurring bombesin by binding to the receptor and, by virtue of the non-peptide bond, failing to exhibit the in vivo activity of naturally occurring bombesin.

EP 0 309 297 A3



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	US-A-4 207 311 (M.R. BROWN et al.) * Claims 1-24 * ---	1	C 07 K 7/00 A 61 K 37/02 C 07 K 7/02
X	NATURE, vol. 316, 29th August 1985, pages 823-826, Mac Millan Journals, Basingstoke, GB; F. CUTTITTA et al.: "Bombesin-like peptides can function in human small-cell lung cancer" * Claims 1-24 * ---	1	
X	AMERICAN JOURNAL OF PHYSIOLOGY, vol. 252, 1987, pages G439 - G442; P. HEINZ-ERIAN et al.: "[D-Phe12]bombesin analogues: a new class of bombesin receptor antagonists" * Page G439, column 1; page G440, figure 1 * ---	1	
P,X	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 263, 1988, pages 5056-5060, Baltimore, US; D.H. COY et al.: "Probing peptide backbone function in bombesin" * Page 5056, column 1; page 5058; page 5059, column 1 * -----	1-17	TECHNICAL FIELDS SEARCHED (Int. Cl.4) C 07 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 06-04-1990	Examiner PEETERS J.C.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			